arXiv:1611.00972v1 [math.RA] 3 Nov 2016

Comments, corrections, and related references welcomed, as always!

TEXed November 4, 2016

STRONG INNER INVERSES IN ENDOMORPHISM RINGS OF VECTOR SPACES

GEORGE M. BERGMAN

Abstract. For V a vector space over a field, or more generally, over a division ring, it is well-known that every $x \in \mathrm{End}(V)$ has an *inner inverse*, i.e., an element $y \in \mathrm{End}(V)$ satisfying $xyx = x$. We show here that a large class of such x have inner inverses y that satisfy with x an infinite family of additional monoid relations, making the monoid generated by x and y what is known as an *inverse monoid* (definition recalled). We obtain consequences of these relations, and related results.

P. Nielsen and J. Šter [13] show that a much larger class of elements x of rings R, including all elements of von Neumann regular rings, have inner inverses satisfying arbitrarily large *finite* subsets of the abovementioned set of relations. But we show by example that the endomorphism ring of any infinite-dimensional vector space contains elements having no inner inverse that simultaneously satisfies all those relations.

A tangential result proved is a condition on an endomap x of a set S that is necessary and sufficient for x to belong to an inverse submonoid of the monoid of all endomaps of S.

1. Background

A basic property of the endomorphism ring $R = \mathrm{End}(V)$ of a vector space V over a division ring is that for every $x \in R$, there exists a $y \in R$ such that

$$xyx \;=\; x. \tag{1}$$

Such a y is called an "inner inverse" to x. (Note that for any endomap x of any set S, an inner inverse to x, i.e., an endomap y of S which satisfies (1), is simply a map that carries every element of the image of x to a preimage of itself under x. For V a vector space and $x \in \mathrm{End}(V)$, an inner inverse to x in $\mathrm{End}(V)$ can be constructed by mapping each member of a vector-space basis B_0 of $x(V)$ to a preimage of itself, and acting arbitrarily on the remaining elements of a basis $B \supseteq B_0$ of V.) If R is a ring and $y \in R$ is an inner inverse of $x \in R$, we see from (1) that xy and yx are idempotent elements of R. A ring such as $\mathrm{End}(V)$ in which every element has an inner inverse is called *von Neumann regular*, often shortened to *regular*.

In general, if y is an inner inverse to x, this does not make x an inner inverse to y. (For instance, every element of a ring is an inner inverse to the element 0.) However, if an element x has an inner inverse y_0, it also has an inner inverse y to which it is, itself, an inner inverse, namely $y = y_0xy_0$.

I had naively thought that $yxy = y$ was the strongest additional relation one could hope to ask of an inner inverse y to an element x in a general regular ring. Hence I was surprised to see, in an unpublished note by Kevin O'Meara, a construction, from any x in such a ring, of an inner inverse y which not only satisfies this additional relation, but also the relation saying that the idempotents xy and yx commute. Note that this commutativity relation, $xyyx = yxxy$, implies that $xxyyxx = x(yxxy)x = (xyx)(xyx) = xx$; in other words, that y^2 is an inner inverse to x^2; and by the symmetry of the relations satisfied, x^2 is also an inner inverse to y^2.

This suggested that one try to obtain an inner inverse y to x that satisfied further relations making y^n an inner inverse to x^n for higher n. And indeed, looking at a particularly nice class of regular rings, namely direct products of matrix rings (in general, of unbounded sizes) over fields, I was able to find an infinite family of relations that an inner inverse to an element of such a ring can be made to satisfy, which includes the relations $x^ny^nx^n = x^n$ and $y^nx^ny^n = y^n$.

A second surprise came when I mentioned this to Pace Nielsen. It turned out that in a paper [13] that he and Janez Šter had submitted for publication, they showed that for any ring R, any element $x \in R$,

2010 *Mathematics Subject Classification.* Primary: 16E50, 16S50, 16U99, 20M18. Secondary: 16S15, 16S36.

Key words and phrases. endomorphism ring of a vector space; inner inverse to a ring element; inverse monoid.

This preprint is readable online at http://math.berkeley.edu/~gbergman/papers/ .

and any positive integer n such that $x,\ x^2, \dots, x^n$ all have inner inverses, one can find an inner inverse y to x satisfying many of the same relations that I had found; in particular, such that y^j is an inner inverse to x^j for all $j \leq n$. After this discussion, they were able to extend the relations they obtained so that the union, over all n, of their families of relations coincided with my family. (Their result appears, so revised, as [13, Theorem 4.8]. Moreover, the form in which they had originally formulated their relations led to improvements in the present note. So [13] and this note each contain material inspired by the other.)

The family of relations in x and y referred to above are monoid relations, and a third surprise was to discover, after drafting most of these pages, that the monoid $\mathcal{F}$ that they define is well known to semigroup theorists, as the free *inverse monoid* on one generator x, and that there is considerable literature on inverse monoids and their monoid algebras, e.g., [14], [8], parts of [5] and [6], [1, §4], [9], and [11].

In §2 below, I develop that monoid essentially "as I found it" (though I have borrowed the notation $\mathcal{F}$ from [1, §4]). In §3, the concept of inverse monoid is sketched, and in §4 we verify the characterization of $\mathcal{F}$ in terms of that concept. In subsequent sections, though $\mathcal{F}$ is referred to in those terms, the results are mostly independent of the literature on the subject.

2. The monoid $\mathcal{F}$

Let us motivate the monoid we will be studying by considering a natural family of pairs of mutually inner inverse vector-space maps. We write maps to the left of their arguments, and compose them accordingly.

Suppose V is an n-dimensional vector space with basis $\{b_1, \dots, b_n\}$, let $x : V \to V$ be the map that sends b_h to b_{h+1} for $h < n$ and sends b_n to 0, and let $y : V \to V$ be the map that sends b_h to b_{h-1} for $h > 1$ and sends b_1 to 0.

Note that for $p, q \geq 0$, the map y^p annihilates the first p of $b_1, \dots, b_n$, while x^q annihilates the last q. (In this motivating sketch, it will help to think of p and q as small compared with n.) If we want to annihilate both the first p and the last q basis elements, we can first apply y^p to get rid of the former, then x^p to bring those that remain back to their original values, and then x^q to annihilate the last q of them; in other words, apply $x^{p+q}y^p$. In addition to annihilating the first p and the last q of the b's, this leaves the b's that remain shifted q steps to the right. If we want to pull them back some distance to the left, we can then apply y^r for some r. Here we may as well take $r \leq p+q$, since if we used a larger value, this would kill some of the elements at the left-hand end of our string, which we could just as well have achieved by using a larger p at the first step. Thus, our combined operation has the form $y^r x^{p+q} y^p$ with $p \geq 0,\ q \geq 0$, and $r \leq p+q$.

Clearly, we could have achieved the same result by *first* annihilating the last q of the b's using x^q, then the first p using y^{p+q}, and finally shifting those that remained using x^{p+q-r}. Thus, $x^{p+q-r}y^{p+q}x^q = y^r x^{p+q} y^p$. Renaming the exponents $p+q-r,\ p+q$ and q as $i,\ j$ and k gives the first set of relations in the next result.

Lemma 1. *The monoid $\mathcal{F}$ presented by two generators x and y, and the infinite system of relations*

$$x^i y^j x^k \ = \ y^{j-i} x^j y^{j-k} \quad \textit{for all } i,\ j,\ k \textit{ such that } 0 \leq i \leq j \textit{ and } 0 \leq k \leq j \tag{2}$$

can also be presented by the subset consisting of the relations

$$x\, y^j x^j \ = \ y^{j-1} x^j, \quad y^j x^j y \ = \ y^j x^{j-1}, \quad \textit{for } 1 \leq j \tag{3}$$

(the cases of (2) where the 3-tuple (i,j,k) has one of the forms $(1,j,j)$ or $(0,j,j-1)$, with the two sides of the latter relation interchanged); and, likewise, by the subset consisting of the relations

$$x^j y^j x \ = \ x^j y^{j-1}, \quad y\, x^j y^j \ = \ x^{j-1} y^j, \quad \textit{for } 1 \leq j \tag{4}$$

(the cases where (i,j,k) has one of the forms $(j,j,1)$, $(j-1,j,0)$, with the same reversal of sides).

In $\mathcal{F}$, every element is equal to the left-hand side of (2) for a unique choice of $i,\ j,\ k$ satisfying the indicated inequalities; hence, equivalently, to the right-hand side of (2) for the same $i,\ j,\ k$.

$\mathcal{F}$ embeds in the direct product of monoids $\langle x, y \mid xy = 1\rangle \times \langle x, y \mid yx = 1\rangle$ via the homomorphism $x \mapsto (x, x),\ y \mapsto (y, y)$.

Proof. Since the set of relations (3) is a subset of (2), to get our first assertion we must show that (3) implies all the relations in (2). Let us first show that it implies the family

$$x^i y^j x^j \ = \ y^{j-i} x^j, \quad y^j x^j y^i \ = \ y^j x^{j-i}, \quad \text{for } 0 \leq i \leq j. \tag{5}$$

Here the cases with $i = 0$ are vacuous, and those with $i = 1$ are (3). For $i > 1$, we note that the left-hand side of the first equation of (5) can be written $x^{i-1}(x\, y^j x^j)$, which by (3) reduces to $x^{i-1} y^{j-1} x^j$.

Writing this as $(x^{i-1}y^{j-1}x^{j-1})\,x$, we may assume by induction on i that this can be reduced using (3) to $(y^{(j-1)-(i-1)}x^{j-1})\,x = y^{j-i}x^j$, the desired expression. The second equation of (5) is obtained by the same calculation, with the order of factors reversed and x and y interchanged.

To get the full set of relations (2), take any i, j, k as in those relations, and note that the expression $x^iy^jx^jy^{j-k}$ can be reduced using (5) in two ways: On the one hand, $x^i\,(y^jx^jy^{j-k}) = x^i\,(y^jx^{j-(j-k)}) = x^iy^jx^k$, which is the left-hand side of (2); on the other hand, $(x^iy^jx^j)y^{j-k} = (y^{j-i}x^j)y^{j-k}$, which is the right-hand side. So these are equal, as desired.

By left-right symmetry, (4) is similarly equivalent to (2).

Let us show next that every $a \in \mathcal{F}$ can be represented by an expression as in (2). Let w be an expression for a of minimal length in x and y. We can write w as an alternating product of nonempty "blocks" of x's and y's. If it consists of ≤ 2 blocks, it is immediate that it has one of the forms shown in (2), so assume it has at least 3 blocks. I claim that w cannot consist of ≥ 4 blocks. For if it does, consider two adjacent blocks which are neither the rightmost nor the leftmost pair. By the right-left symmetry of the statements we are dealing with, we may assume for simplicity that the length of the first of these blocks is at least that of the second, and by symmetry in x and y, that the first block is a power of x and the second a power of y; so our product of two blocks can be written x^iy^j with $i \geq j \geq 1$. Since this pair of blocks does not occur at the right end of w, it must be followed by an x; so w has a subword x^iy^jx, which in turn has the subword x^jy^jx, which can be reduced by (4) to the shorter word x^jy^{j-1}, contradicting our minimality assumption. So w must have just three blocks, and we can assume without loss of generality that it has the form $x^iy^jx^k$. If j is $\geq$ both i and k, we are done. If not, assume without loss of generality that $i > j$. Then w has the subword x^jy^jx, which, as above, contradicts minimality. Putting aside our "w.l.o.g" assumptions, what we have shown is that a word w of minimal length representing a will be of one of the forms shown in (2). Whichever of those forms it has, (2) allows us to represent a in the other form as well (though that may not be a minimal-length expression for it).

To show that the expression for a in each of these forms is unique, and simultaneously obtain the embedding of the final assertion of our lemma, let us first note that the generators x and y of each of the monoids $\langle x, y \mid xy = 1\rangle$ and $\langle x, y \mid yx = 1\rangle$ clearly satisfy (3). Hence we get a homomorphism

$$h : \mathcal{F} \;\to\; \langle x, y \mid xy = 1\rangle \;\times\; \langle x, y \mid yx = 1\rangle \tag{6}$$

carrying x to (x, x) and y to (y, y). Now in $\langle x, y \mid xy = 1\rangle$, the expressions y^ix^j give a normal form (as is easily checked by the method of [3]), while in $\langle x, y \mid yx = 1\rangle$, the same is true of the expressions x^iy^j. The image under h of an element $x^iy^jx^k$ with $0 \leq i \leq j$ and $0 \leq k \leq j$, written using these normal forms, is $(y^{j-i}x^k, x^iy^{j-k})$. Here the first component determines k, the second determines i, and with these known, either component determines j. So distinct expressions $x^iy^jx^k$ as in (2) have distinct images under h, showing, on the one hand, that (6) is an embedding, and on the other, that distinct expressions as in the left-hand side of (2) represent distinct elements of $\mathcal{F}$. □

Intuitively, the first relation of (3) says that after a factor x^jy^{j-1}, one can "drop" a factor yx; in other words, that x^jy^{j-1} behaves like x in that respect. The other relation of (3) and those of (4) have the obvious analogous interpretations. More generally, I claim that

(7) If w is a word in x and y which has strictly more x's than y's, then in the monoid $\mathcal{F}$ we have $wyx = w = xyw$.

(8) If w is a word in x and y which has strictly more y's than x's, then in the monoid $\mathcal{F}$ we have $wxy = w = yxw$.

Indeed, by symmetry in x and y, it suffices to prove (7), and by right-left symmetry it suffices to prove the first relation thereof, $wyx = w$. The last sentence of Lemma 1 shows that it suffices to prove that relation in each of the monoids $\langle x, y \mid xy = 1\rangle$ and $\langle x, y \mid yx = 1\rangle$. The result is trivially true in the latter monoid. In the former, w can be reduced to a word y^ix^j such that $j - i = (\text{number of } x\text{'s in } w) - (\text{number of } y\text{'s in } w)$. This difference is positive by the hypothesis of (7), so $j > 0$; so w is equal in $\langle x, y \mid xy = 1\rangle$ to a word ending in x, so by the relation $xyx = x$ we indeed have $wyx = w$.

3. A quick introduction to inverse monoids

The concept of *inverse monoid*, mentioned in the Introduction, is motivated by the natural structure on the set of *partial one-to-one maps* of a set S into itself; i.e., the one-to-one maps from subsets of S to S.

Clearly, the composite of two partial one-to-one maps is again such a map, as is the identity map of S, so such maps form a monoid. Further, if x is such a partial map, and we view it as a relation, hence a subset of $S \times S$, then the inverse relation $y = \{(t,s) \mid (s,t) \in x\}$ is again a partial one-to-one map. If we consider all identities satisfied by such partial maps, under the monoid operations and this "inverse" operation, these determine a *variety* of monoids with an additional unary operation. We shall call objects of this variety *inverse monoids*.

Among semigroup-theorists, the "inverse" operation is generally written $y = x^{-1}$. But this would conflict badly with the usage of ring theory, so, following [1, §4], I will instead write x^*.

It turns out that (as with the inverse operation of a group) if a monoid admits any unary operation * satisfying the identities of the above variety, then that operation is unique [5, Theorem 1.17], [8, Theorem 1.3]; so (like groups) inverse monoids can be identified with a subclass of the monoids, and they are in fact generally so described in the literature. Precisely, it is easy to see that the identities of inverse monoids include $x = xx^*x$ and $x^* = x^*xx^*$, and the standard definition is that an inverse monoid is a monoid M such that

(9) For every $x \in M$ there is a *unique* $y \in M$ satisfying $xyx = x$ and $yxy = y$ [8, p. 6, conditions (1) and (2)], [14, Definition II.1.1], [5, p. 28, lines 9-11].

(Caveat: Given an element x in an inverse monoid, an element y satisfying $xyx = x$, but not necessarily $yxy = y$, will not, in general, be unique. For instance, in the monoid of partial one-to-one endomaps of a set S, if x is the empty partial map, every y satisfies that relation.)

We remark that most of the literature describes its subject as *inverse semigroups,* where no identity element is assumed. Those that are monoids are indeed considered, but as a subcase. However, as with the relationship between ordinary semigroups and monoids, or nonunital and unital rings, either version of the theory can be reduced to the other, so which one treats as primary is largely a matter of taste; and I will talk about *inverse monoids* here.

The most commonly used characterizations of inverse monoids are not given by identities; (9) is such a characterization. We recall two others: if M is a monoid and * a unary operation on its underlying set, then each of the following conditions is equivalent to $(M,^*)$ being an inverse monoid.

(10) $(M,^*)$ can be embedded in the monoid of one-to-one endomaps of some set S, in such a way that * acts as the relational inverse map [14, Corollary IV.1.9], [8, p. 36, Theorem 1.5.1], [5, Theorem 1.2].

(11) For every $x \in M$, one has $x = xx^*x$, and, further, every pair of idempotent elements of M commute [8, p. 6, Theorem 3], [5, Theorem 1.17(i)], cf. [14, Theorem II.1.2].

To move toward a characterization by identities, note that the identities satisfied by partial one-to-one operations on a set include the conditions making * an *involution* of M,

$$(ab)^* = b^*a^* \quad \text{and} \quad a^{**} = a. \tag{12}$$

From these it follows that if an inverse monoid $(M,^*)$ is generated by a subset X, then as a monoid, M is generated by $X \cup X^*$. Indeed, the submonoid generated by that set will contain X and be closed under the monoid operations, and by (12) it will be closed under *, hence it will be all of M.

Now the reason why (11) is not a set of identities is that the condition that idempotents commute is not an equation in arbitrary elements of M. However, given the identity $x = xx^*x$ of (11), we see that if e is an idempotent, then $e = ee^*e = e(e^*e^*)e = (ee^*)(e^*e^{**})$, a product of two idempotents of the form xx^*. Hence if we combine the monoid identities, the identity $x = xx^*x$, and the identity saying that any two elements of the forms xx^* and yy^* commute, then these together imply (11), and so define the variety of inverse monoids.

A feature that inverse monoids share with groups, though not nearly as easy to prove, is

(13) If $(M,^*)$ is an inverse monoid, and N is a homomorphic image of M as a monoid, then the operation * on M induces an operation on N, whence N becomes an inverse monoid [6, Theorem 7.36], [14, Lemma II.1.10].

In view of the uniqueness of the inverse monoid structure on a monoid when one exists, I will sometimes take the shortcut of saying that a monoid "is" an inverse monoid (as in the standard usage in the field) when I mean that it admits an operation * making it an inverse monoid.

In addition to groups, and monoids of partial one-to-one maps of sets, another important class of inverse semigroups are the semilattices, where * is taken to be the identity map.

For extensive developments of the theory of inverse semigroups, see [8] and [14].

4. $\mathcal{F}$ as a free inverse monoid

In §2, we motivated the structure of our monoid $\mathcal{F}$ in terms of certain endomorphisms of finite-dimensional vector spaces. What we did can be looked at as a partial-one-one-map construction in a different guise. Each of x and y, and hence also every monoid word in those elements, sends certain members of our basis $\{b_1, \ldots, b_n\}$ to members of that basis, and does so in a one-to-one fashion, while it maps the remaining elements of that basis to 0. If we associate to every endomorphism z of V that acts in such a way on our basis the partial endomap of that basis which agrees with z on elements that z sends to basis elements, and is undefined on those z sends 0, then the resulting partial maps compose like the corresponding linear maps.

On each of our finite-dimensional vector spaces, the monoid of endomorphisms generated by x and y is finite; but if we think of x and y as acting simultaneously on n-dimensional spaces for all natural numbers n, we shall find that we get the infinitely many distinct elements of $\mathcal{F}$. We will deal with actions on vector spaces in the next section; here, let us show monoid-theoretically that

Lemma 2. *The monoid $\mathcal{F}$ of Lemma 1 is the free inverse monoid on one generator x, with $y = x^*$.*

Proof. To see that $\mathcal{F}$ is an inverse monoid, we shall use its representation, established in Lemma 1, as a subdirect product of

$$\langle x, y \mid xy = 1 \rangle \quad \text{and} \quad \langle x, y \mid yx = 1 \rangle. \tag{14}$$

These are two copies of the same monoid, called the *bicyclic* monoid, which has a natural representation by partial endomaps of the natural numbers, with the left-invertible generator acting as the right shift and the right-invertible generator as the left shift (undefined at 0). The involution * of each of the inverse monoids of (14) interchanges the generators x and y of that monoid, so the submonoid $\mathcal{F}$ generated by (x, x) and (y, y) is closed under coordinatewise application of *, hence is itself an inverse monoid.

To show that the inverse monoid $(\mathcal{F}, ^*)$ is free on $\{x\}$, it suffices to show that for any element x of an inverse monoid, if we write $x^* = y$, then x and y satisfy the relations of (3). To get the first of these relations, let us write xy^jx^j as $(xy)(y^{j-1}x^{j-1})x$. Since $y = x^*$, we see from (12) that $(y^{j-1})^* = x^{j-1}$, hence both xy and $y^{j-1}x^{j-1}$ are idempotent, hence by (11) they commute, so we can write $(xy)(y^{j-1}x^{j-1})x$ as $(y^{j-1}x^{j-1})(xy)x = y^{j-1}x^{j-1}(xyx) = y^{j-1}x^{j-1}x = y^{j-1}x^j$, giving the desired relation. Applying the involution *, we get the other relation of (3). □

The normal form for elements of the free inverse monoid on one generator given by the expressions on either side of (2) is also obtained in [1, proof of Lemma 4.1]. I do not know whether a system of defining monoid relations as economical as (3) or (4) has previously been noted.

The multiplicative monoids of the rings we will be looking at in this note are not, in general, inverse monoids; but we will nonetheless be interested in pairs of elements x and y of these rings that satisfy the relations holding between elements x and x^* of an inverse monoid. Let us therefore make

Definition 3. *If x is an element of a monoid M, we shall call an element $y \in M$ a* strong inner inverse *to x if the submonoid of M generated by x and y can be made an inverse monoid with $y = x^*$; equivalently, if there exists a monoid homomorphism $\mathcal{F} \to M$ carrying $x, y \in \mathcal{F}$ to the elements of M denoted by these same symbols; equivalently, if $x, y \in M$ satisfy any of the equivalent systems of monoid relations* (2), (3), (4).

Clearly, the relation "is a strong inner inverse of" is symmetric.

From our observations on the bicyclic monoid $\langle x, y \mid xy = 1 \rangle \cong \langle x, y \mid yx = 1 \rangle$, we see that any right or left inverse of an element x of a monoid (and hence in particular, a 2-sided inverse) is a strong inner inverse to x.

If every element of a monoid M has a *unique* strong inner inverse, then M will, *a fortiori*, satisfy (9), and so be an inverse monoid. But we shall see below that in many rings, every element has a strong inner inverse, but these are not unique.

(We remark that our use of "strong" in "strong inner inverse" should not be confused with the use of "strongly" in the existing concept of a "strongly regular ring", a regular ring in which all idempotents are central. That condition is much more restrictive than the condition that every element have a strong inner inverse in our sense.)

5. Many vector space endomorphisms have strong inner inverses

We shall show here that if V is a vector space over a division ring D, then a large class of elements $x \in \mathrm{End}_D(V)$ have strong inner inverses. Our proof will make use of the following weak version of Jordan canonical form, which holds over division rings.

Lemma 4. *Suppose V is a finite-dimensional vector space over a division ring D, and x a vector-space endomorphism of V. Let us call a nonzero x-invariant subspace W of V "x-basic" if it admits a basis $b_1, \dots, b_n$ such that $x(b_h) = b_{h+1}$ for $h < n$, and $x(b_n) = 0$. Then V can be written as the direct sum of a subspace on which x acts invertibly, and a family of x-basic subspaces.*

Proof. By Fitting's Lemma [10, Theorem 19.16], there exists an $N \geq 1$ such that

$$V \;=\; \mathrm{im}(x^N) \;\oplus\; \ker(x^N). \tag{15}$$

Clearly, each of these summands is x-invariant, and x acts injectively (hence invertibly) on the former, since otherwise the summands would have nonzero intersection. So it will suffice to show that $\ker(x^N)$ is a direct sum of x-basic subspaces. This can be done in a well-known manner which we now sketch.

Noting that $\ker(x) \cap \mathrm{im}(x^N) \subseteq \ker(x^N) \cap \mathrm{im}(x^N) = \{0\}$, we look at the chain of subspaces

$$\ker(x) \cap \mathrm{im}(x^{N-1}) \;\subseteq\; \ker(x) \cap \mathrm{im}(x^{N-2}) \;\subseteq\; \dots \;\subseteq\; \ker(x) \cap \mathrm{im}(x) \;\subseteq\; \ker(x). \tag{16}$$

We take a basis B_{N-1} for the first of these, extend it to a basis $B_{N-1} \cup B_{N-2}$ for the second, and so on, getting a basis $B_{N-1} \cup \dots \cup B_1 \cup B_0$ for $\ker(x)$, with the B_i disjoint. For each i we now write each $b \in B_i$ as $x^i(b')$ for some $b' \in \ker(x^N)$. If we consider the set

$$\textstyle\bigcup_{0 \leq i \leq N-1,\, b \in B_i} \{x^h(b') \mid\; 0 \leq h \leq i\}, \tag{17}$$

then this is easily shown to form a D-basis for $\ker(x^N)$, and for each $i \leq N-1$ and $b \in B_i$, the span of the set $\{x^h(b') \mid\; 0 \leq h \leq i\}$ is x-basic, since x carries $x^h(b')$ to $x^{h+1}(b')$ for $h \leq i$, and $x^i(b') = b \in B_i \subseteq \ker(x)$ to 0. □

We can now prove

Theorem 5. *Let D be a division ring, V a right vector space over D, and x an endomorphism of V such that V can be written as a direct sum of x-invariant D-subspaces, each of which is either finite-dimensional, or has the property that the action of x on it is one-to-one, or has the property that the action of x on it is surjective. Then x has a strong inner inverse y in $\mathrm{End}_D(V)$, which carries each of those direct summands into itself.*

If, moreover, our decomposition of V includes, for each positive integer n, at least one n-dimensional summand which is x-basic in the sense of Lemma 4, then a strong inner inverse y to x can be chosen so that the ring of additive-group endomorphisms of V generated by the action of D and the actions of x and y has precisely the structure of the monoid ring $D\mathcal{F}$. In particular, if D is a field k, this says that the k-algebra of endomorphisms of V generated by x and y is isomorphic to the monoid algebra $k\mathcal{F}$.

Proof. In proving the first assertion we may, by the preceding lemma, assume that each of the given x-invariant summands of V either has x acting surjectively on it, or has x acting injectively on it, or is x-basic; so we are reduced to proving that x has a strong inner inverse if x acts in one of these three ways on V itself.

If x is surjective or injective, then it is right or left invertible in $\mathrm{End}(V)$, and a right or left inverse will be the desired strong inner inverse.

So suppose V is x-basic, and let $\{b_1, \dots, b_n\}$ be a basis of V such that x carries each b_h with $h < n$ to b_{h+1}, and carries b_n to 0. As in the motivating sketch of §2, define y to carry b_h to b_{h-1} for $h > 1$, and b_1 to 0. We could verify that y is a strong inner inverse to x using the correspondence between one-to-one partial maps on our basis of V and certain endomorphisms of V, as noted at the beginning of §4, but the hands-on proof that x and y satisfy the relations of (3) is quick enough, so I will sketch it.

To prove, first, the relation $x\,y^j x^j = y^{j-1} x^j$ for $1 \leq j$, let us compare the effect of both sides on some b_h. If $j + h > n$, then the x^j at the right end of each monomial annihilates b_h, so the two sides of the relation indeed agree on b_h. If $j + h \leq n$ (so in particular, $h < n$), it is immediate to check that each side gives b_{h+1}, and the relation again holds.

Similarly, on checking the results of applying the two sides of the relation $y^j x^j y = y^j x^{j-1}$ for $1 \leq j$ to b_h, we find that if $h = 1$ or $h + (j-1) > n$, both sides give 0, while in the contrary case, both sides give b_{h-1}. This completes the proof of the first assertion of the theorem.

To get the second assertion, let y be constructed on each summand of our decomposition of V as in the proof of the first assertion. Since y is a strong inner inverse to x, the elements x and y satisfy the relations defining $\mathcal{F}$, and since they are D-linear, they commute with the action of the elements of D; hence the actions of x, y and the elements of D yield a $D\mathcal{F}$-module structure on V. It remains to show that this module is faithful, so let $a \in D\mathcal{F} - \{0\}$, and let us show that the action of a on V is nonzero.

To this end, let $x^i y^j x^k$, with i, j, k as in (2), be the element of $\mathcal{F}$ having nonzero D-coefficient in a which gives the *least* 3-tuple (k, j, i) under lexicographic order. (Note the reversed order of indices.) Choose an x-basic subspace V' in our given decomposition of V which has dimension exactly $j+1$, and let $\{b_1, \dots, b_{j+1}\}$ be a basis for V' on which x and y act as right and left shift operators.

I claim that the element b_{j-k+1} is not annihilated by a. Note first how $x^i y^j x^k$ acts on b_{j-k+1}. The factor x^k carries it to b_{j+1} (note that any higher power of x would kill it), y^j carries this to b_1 (here any higher power of y would kill it), and x^i brings this to b_{i+1}. So it will suffice to show that none of the other terms occurring in a carry b_{j-k+1} to an expression in which b_{i+1} appears. Let $x^{i'} y^{j'} x^{k'}$ be any other term occurring in a.

By our minimality assumption on (k, j, i), we must have $k' \geq k$. If this inequality is strict, we see that b_{j-k+1} is killed on applying $x^{k'}$ to it. On the other hand, if $k' = k$, then $j' \geq j$, and if this inequality is strict, our basis element is killed on applying $y^{j'} x^k$. Finally, if $k' = k$ and $j' = j$, we must have $i' \neq i$, and we see that $x^{i'} y^j x^k$ will carry b_{j-k+1} to $b_{i'+1} \neq b_{i+1}$. Hence in $a(b_{j-k+1})$, only the term $x^i y^j x^k (b_{j-k+1})$ contributes to the coefficient of b_{i+1}, so $a(b_{j-k+1}) \neq 0$, so a indeed acts nontrivially on V. □

We shall see in the next section that in the first assertion of the above theorem, the restriction to endomorphisms x such that V is an appropriate direct sum of sorts of x-invariant subspaces cannot be dropped. Here is a case where that condition holds automatically.

Corollary 6. *Let D be a division ring, and R the direct product of the endomorphism rings of a family of finite-dimensional D-vector-spaces, $R = \prod_{i \in I} \mathrm{End}(V_i)$ (i.e., a direct product of full matrix rings of various sizes over D). Then every element of R has a strong inner inverse under multiplication.*

Proof. Apply the preceding theorem with $V = \bigoplus V_i$, and R identified with the ring of endomorphisms of V that carry each V_i into itself. □

We mentioned earlier that for M a monoid, the statement that every element of M has a strong inner inverse does not entail that such strong inner inverses are unique. For an explicit example, consider the monoid $\mathrm{End}(V)$, where V is a vector space with basis $\{b_1, \dots, b_n\}$ for some $n > 1$, and x and y again act by right and left shifts. Then x and y are nilpotent, and do not commute. Since x is nilpotent, $1 + x$ is invertible, hence conjugation by that element is an automorphism of $\mathrm{End}(V_i)$ which fixes x but not y. Hence $(1+x)^{-1} y (1+x)$ is a strong inner inverse to x distinct from y.

One may ask

Question 7. *Given endomorphisms $x_1, \dots, x_r$ of a vector space V over a field, or more generally, over a division ring, under what natural conditions can we find strong inner inverses $y_1, \dots, y_r$ to these elements, such that the monoid generated by $x_1, \dots, x_r$ and $y_1, \dots, y_r$ is an inverse monoid?*

If $r > 1$ such strong inner inverses need not exist for general $x_1, \dots, x_r \in \mathrm{End}(V)$, even if V is finite-dimensional. For instance, if we take x_1 and x_2 to be noncommuting idempotents, no submonoid of $\mathrm{End}(V)$ containing them satisfies (11). A sufficient condition is that there exist a basis B for V such that every x_i carries each member of B either to another member of B, or to 0, and acts in a one-to-one fashion on those that it does not send to 0. But this is not necessary. For example, if we take for $x_1, \dots, x_r$ any automorphisms of V, then they generate a group, which is an inverse monoid; but if any of them has determinant $\neq \pm 1$, it cannot permute a basis of V.

(I will mention one result that has a vaguely related feel. Given a vector space V and a finite family of subspaces $S_1, \dots, S_n$, one may ask under what conditions there exists a basis B for V such that each S_i is the subspace spanned by a subset $B_i \subseteq B$. By [4, Exercise 6.1:16], such a basis exists if and only if the lattice of subspaces generated by $S_1, \dots, S_n$, is distributive.)

There is an interesting analog, for families of endomorphisms $x_1, \dots, x_r$ of vector spaces, of the class of x-basic spaces. Namely, for every finite connected subgraph S of the Cayley graph of the free group on r generators $g_1, \dots, g_r$, let V_S be a vector space with a basis $\{b_s\}$ indexed by the vertices s of S, and for $i = 1, \dots, r$, let x_i act on V_S by taking b_s to b_t if S has a directed edge from s to t indexed by g_i, to

0 if no edge indexed by g_i comes out of s; and likewise, let x_i^* act by taking b_t back to b_s in cases of the first sort, while taking b_t to 0 if t has no edge indexed by g_i coming into it. From a description of the free inverse monoid on r generators due to Munn [12] [14, §VIII.3], [8, §6.4], one finds that the algebra of operations on $\bigoplus_S V_S$ generated by the resulting maps x_i and x_i^* is isomorphic to the monoid algebra of the free inverse monoid on r generators.

6. But others do not

Let us show that in Theorem 5, the hypothesis that V have a decomposition as a direct sum of well-behaved x-invariant subspaces cannot be dropped. We will need the next result, which concerns monoids of ordinary (everywhere defined, not necessarily one-to-one) endomaps of sets.

Lemma 8. *Let x be an endomap of a set S. Then a necessary condition for x to have a strong inner inverse in the monoid of all endomaps of S is*

$$x\,(\textstyle\bigcap_{n\geq 0} x^n(S)) \;=\; \bigcap_{n\geq 0} x^n(S). \tag{18}$$

Proof. Suppose x has a strong inner inverse y. Since in (18), the relation "$\subseteq$" clearly holds, we must show the reverse inclusion.

So consider any $s \in \bigcap x^n(S)$. In view of the relations $x^n = x^n y^n x^n$, we have

$$s \;=\; x^n y^n s \quad \text{for all } n \geq 0. \tag{19}$$

If we take $n \geq 1$, apply y to both sides of this equation, and then apply to the right-hand side the second relation of (4) with n for j, we get $ys = x^{n-1}y^n s$. Hence $ys \in \bigcap x^n(S)$; and applying x to both sides of this relation, and invoking the $n = 1$ case of (19), we conclude that $s \in x\,(\bigcap x^n(S))$, as desired. □

We shall see in the next section that the condition of the above lemma is sufficient as well as necessary; but we only need necessity for the example below.

The desired vector-space example is obtained by slightly tweaking the well-behaved example, implicit in Theorem 5, of a space V which is a direct sum of x-basic subspaces of all natural number dimensions.

Proposition 9. *Let V be a vector space over a division ring D, with a basis consisting of elements $b_{n,i}$ for all positive integers n and i with $i \leq n$, and one more basis element, b_+; and let $x \in \mathrm{End}(V)$ be given by*

$$x(b_{n,i}) = b_{n,i+1} \text{ if } i < n, \quad x(b_{n,n}) = b_+ \text{ for all } n, \quad x(b_+) = 0. \tag{20}$$

Then x has no strong inner inverse $y \in \mathrm{End}(V)$ (in fact, no strong inner inverse in the monoid of all set-maps $V \to V$).

Proof. Clearly, $\bigcap_{n\geq 0} x^n(V)$ is the 1-dimensional subspace of V spanned by b_+. The image of this subspace under x is the zero subspace, so (18) is not satisfied, hence Lemma 8 gives the desired conclusion. □

The next result gets some further mileage out of the above example. The first assertion of that proposition answers a question posed in an earlier version of [13]; the second shows that in Lemma 8 and Proposition 9, the relations characterizing a strong inner inverse y to x cannot be replaced by the subset consisting of the relations $x^n y^n x^n = x^n$ and $y^n x^n y^n = y^n$ for all $n \geq 0$.

Proposition 10. *Let V and x be as in Proposition 9. Let V_n, for each $n \geq 1$, be the subspace of V spanned by $b_{n,1}, \ldots, b_{n,n}$, let V_+ be the subspace of V spanned by b_+, let R_0 be the ring of endomorphisms of V carrying each of these subspaces into itself, and let L be the space of endomorphisms of V of finite rank.*

Then $R = R_0 + L$ is a unit regular ring *containing x, but containing no y that satisfies $x^n y^n x^n = x^n$ for all n.*

On the other hand, the full ring $\mathrm{End}(V)$ *contains an element y which satisfies both $x^n y^n x^n = x^n$ and $y^n x^n y^n = y^n$ for all $n \geq 0$.*

Proof. Let us prove the above claims in reverse order: first (easiest) the existence of a y as in the final assertion, then the non-existence result of the first assertion, and finally, the unit-regularity of $R = R_0 + L$.

A y as in the final assertion is defined by the familiar formulas $y(b_{n,i}) = b_{n,i-1}$ for $i > 1$, together with the unexpected formulas $y(b_+) = b_{1,1}$ and $y(b_{n,1}) = b_{n+1,1}$ $(n \geq 1)$. In checking that for every n we have $x^n y^n x^n = x^n$, let us think of that relation as saying that $x^n y^n$ fixes all elements of $x^n(V)$. Now $x^n(V)$ is spanned by the elements $b_{m,i}$ with $i > n$, and b_+; and it is straightforward to check that elements of

each of these sorts are fixed by $x^n y^n$. Similarly, the desired relation $y^n x^n y^n = y^n$ says that all elements of $y^n(V)$ are fixed by $y^n x^n$. Now when y is applied to a basis element $b_{m,i}$, we see that the difference $m - i$ is increased by 1, whether we are in the case $i > 1$ or $i = 1$; so the result of applying y^n to a basis element $b_{m,i}$ is a basis element of the form $b_{m',i'}$ with $m' - i' \geq n$. Again, it is immediate to check that any basis element with this property is fixed by $y^n x^n$. On the other hand, in evaluating $y^n x^n y^n(b_+)$, we can use the fact that $b_+ \in x^n(V)$, so by our observations on the relations $x^n y^n x^n = x^n$, b_+ is fixed under $x^n y^n$. Now applying y^n, we get $y^n x^n y^n(b_+) = y^n(b_+)$, as desired.

We turn next to $R = R_0 + L$. It is clear that R is a ring and L an ideal of R. The endomorphism of V that carries all basis elements of the form $b_{n,n}$ to b_+, and all other basis elements to zero, has rank 1, hence belongs to L, and if we subtract it from x we get a member of R_0; so $x \in R$.

To prove the nonexistence of a $y \in R$ satisfying the relations $x^n y^n x^n = x^n$, consider any $y = y_0 + y_1 \in R$, where $y_0 \in R_0$ and $y_1 \in L$. Since y_1 has finite rank, its range lies in the sum of V_+ and finitely many of the V_n. Also, by the description of V, the element $y(b_+)$ lies in the sum of V_+ and finitely many of the V_n. Hence we can choose an $n_0 > 0$ such that both the space $y_1(V)$ and the element $y(b_+)$ have zero projection in all of the spaces V_n for $n \geq n_0$.

Now consider the element $x^{n_0} y^{n_0} x^{n_0}(b_{n_0,1})$. The x^{n_0} that acts on $b_{n_0,1}$ carries it to b_+, and subsequent iterations of y will, by our choice of n_0, carry this into a member of $V_+ + \sum_{n<n_0} V_n$. But every member of this sum is annihilated by x^{n_0}; so $x^{n_0} y^{n_0} x^{n_0}(b_{n_0,1}) = 0$, though $x^{n_0}(b_{n_0,1}) = b_+$. Hence $x^{n_0} y^{n_0} x^{n_0} \neq x^{n_0}$, as claimed.

It remains to show that R is unit regular; i.e., that every element has an inner inverse which is a unit. We shall recall sufficient conditions for this to hold, given in [2, Lemma 3.5, p. 600] for a general ring R with an ideal L, in terms of properties of R, R/L and L, and verify that these hold in the case at hand. (I am indebted to Ken Goodearl and Pace Nielsen for supplying the tools for this part of the proof.)

First, we must know that R is regular. This follows from [7, Lemma 1.3], since L, being an ideal of $\mathrm{End}(V)$, is regular, and $R/L \cong R_0/(R_0 \cap L)$, a homomorphic image of the regular ring R_0, is also regular.

Next, we must know that eLe is unit regular for every idempotent $e \in L$. But $eLe \cong \mathrm{End}(eV)$, the endomorphism ring of a finite dimensional vector space, and so is unit regular by [7, Theorem 4.1(a) $\iff$ (c)].

We must also know that R/L is unit regular. For this we again use the fact that $R/L \cong R_0/(R_0 \cap L)$, this time combined with the fact that R_0, a direct product of unit regular rings, is unit regular.

Finally, we need to know that every unit of R/L lifts to a unit of R. Now a unit of R/L and its inverse will be images of elements $u, v \in R_0$. Hence $uv-1$ and $vu-1$ will both lie in $R_0 \cap L$, hence will be members of R_0 whose components equal zero in all but finitely many of the algebras $\mathrm{End}(V_n)$ and $\mathrm{End}(V_+)$. Hence if we modify u and v only in their behaviors on the finitely many factors where they are not inverse to one another, replacing their actions there with, say, the identity elements of those factors, we get elements $u', v' \in R_0$ which *are* inverse to one another, and these give the desired liftings to R of the original unit and its inverse in R/L. □

In contrast to the first assertion of the above proposition, we know from [13, Theorem 4.8], mentioned in the Introduction, that the x of the above example has inner inverses in R which satisfy any *finite* subset of the relations (2).

7. Digression: Characterization of set-maps having strong inner inverses

Here is the promised strengthening of Lemma 8. (It will not be used in subsequent sections.)

Theorem 11. *Let x be an endomap of a set S. Then condition* (18) *is necessary and sufficient for x to have a strong inner inverse y in the monoid of all endomaps of S.*

Proof. In view of Lemma 8, we only have to prove sufficiency, so assume x satisfies (18), and let us construct y.

For every $s \in S$, we define its "depth",

(21) $d(s)$ = greatest integer $n \geq 0$ such that $s \in x^n(S)$ if this exists, or ∞ if $s \in \bigcap_{n \geq 0} x^n(S)$.
(In statements such as (22) and (23) below, we will understand $\infty + 1 = \infty = \infty - 1$.)

In particular, $d(s) = 0$ if and only if $s \notin x(S)$. Clearly,

(22) For all $s \in S$, we have $d(x(s)) \geq d(s) + 1$.

Moreover,

(23) If $d(s) > 0$, then s can be written as $x(t)$ for some t with $d(t) = d(s) - 1$.

Indeed, for $0 < d(s) < \infty$, (23) is straightforward, while the case where $d(s) = \infty$ is our assumption (18).

We will begin the construction of y by defining it on all $s \in x(S)$. The criterion for choosing $y(s)$ will be fairly natural, but with one nonobvious restriction, applying to elements $s \in S$ that satisfy

(24) $d(x^n(s)) \;=\; d(s) + n \quad$ for all $n \geq 0$ (cf. (22)).

For such an s, *if* the function y that we shall define below satisfies

(25) $y\,x^n(s) \;=\; x^{n-1}(s) \quad$ for all $n \geq 1$,

let us call s *y-stable*. We now specify y on $x(S)$ by the rule

(26) For each $s \in x(S)$, let $y(s)$ be an element t such that $s = x(t)$ and $d(t) = d(s) - 1$ (as in (23)). Moreover, if there are any $s \in S$ satisfying (24), then we make our choice of y satisfy (25) for *at least one* such s (y-stable).

Can we achieve the second condition of (26)? If x is one-to-one on the x-orbit of some s satisfying (24), we can clearly define y on elements $x^n(s)$ by (25). If, on the other hand, x is not one-to-one on the orbit of s, then that orbit must eventually become periodic, and taking a new s in the periodic part, x will be one-to-one on the orbit of that element, and we can define y on that orbit by (25). So the second sentence of (26) can indeed be achieved.

Note that, in view of (22), and the condition $d(t) = d(s) - 1$ in (26), we can say that

(27) Once y has been defined on $x(S)$ so as to satisfy (26), we can use this partial definition of y to evaluate $y^n(s)$ for any $s \in S$ with $d(s) \geq n$.

We now want to define y on elements $s \notin x(S)$.

A fairly easy case is that in which d makes a "jump" somewhere on the x-orbit of s:

(28) If $s \in S$ with $d(s) = 0$, and if for some positive integer n, $d(x^n(s)) > n$, then letting n be the least value for which this is true, we define $y(s) = y^{n+1}x^n(s)$.

In view of the hypothesis of (28) that $d(x^n(s)) > n$, we see by (27) that $y^{n+1}x^n(s)$ is determined by the partial definition of y that we have so far; so (28) makes sense.

Finally, let us define y on those $s \notin x(S)$ such that $d(x^n(s)) = n$ for all $n \geq 0$. We will need the following equivalence relation.

(29) For $s, s' \in S$ such that $d(x^n(s)) = n = d(x^n(s'))$ for all $n \geq 0$, we shall write $s \approx s'$ if there exists some n such that $x^n(s) = x^n(s')$.

We now specify

(30) If $s \in S$ satisfies $d(x^n(s)) = n$ for all $n \geq 0$, then we take $y(s)$ to be any y-stable element (any element satisfying (25)), subject only to the condition that if $s \approx s'$, then $y(s) = y(s')$.

To see that we can do this, note that the situation "$d(x^n(s)) = n$ for all $n \geq 0$" is a case of (24), hence by the second sentence of (26), if there exist s as in (30), then there also exist y-stable elements, which is all we need to carry out (30).

This completes the construction of y. Let us record two immediate consequences of the first sentence of (26). First, the fact that $y(s)$ is chosen to be a preimage of s under x can be reworded:

(31) If $d(s) > 0$, then $xy(s) = s$.

Second, combining the condition on $d(t)$ in (26) with the fact that for $s \notin xS$, $d(y(s)) \geq 0 > d(s) - 1$, we see that

(32) For all $s \in S$, we have $d(y(s)) \geq d(s) - 1$ (cf. (22)).

Let us now show that x and the y we have constructed satisfy the relations (3). The first of these, $xy^jx^j = y^{j-1}x^j$, says that for every $s \in S$, the element $y^{j-1}x^j(s)$ is fixed under xy. This is easy: by (22) and (32), every $s \in S$ satisfies $d(y^{j-1}x^j(s)) \geq 1$, so by (31), $y^{j-1}x^j(s)$ is indeed fixed under xy.

The other relation from (3),

(33) $y^jx^jy \;=\; y^jx^{j-1}$,

automatically holds when applied to elements s with $d(s) > 0$, since the two sides differ by a right factor of xy, again allowing us to apply (31).

So suppose $d(s) = 0$.

Assume first that s has the property that $d(x^n(s)) > n$ for some n, and as in (28), let $n \geq 1$ be the least such value. Applying the two sides of (33) to s, and using (28) on the left-hand side, we see that we need to prove

$$y^j x^j y^{n+1} x^n(s) \;=\; y^j x^{j-1}(s). \tag{34}$$

Now because we have assumed the element s satisfies $d(x^n(s)) \geq n+1$, we see that on the left-hand side of (34), each of the terms in the string y^{n+1} gets applied to an element of positive depth; hence by (31), we can repeatedly cancel sequences $x\,y$ at the interface between this and the string x^j. The outcome of these cancellations depends on the relation between j and n in (34). Suppose first that $j > n$. Then the above repeated cancellations turn the left-hand side of (34) into $y^j x^{j-n-1} x^n(s)$, which equals the right-hand, as desired.

If, rather, $j \leq n$, then the cancellation mentioned turns the left-hand side of (34) into $y^j y^{n+1-j} x^n(s) = y^{n+1} x^n(s)$, which by (28) is the value we have assigned to $y(s)$. We want to compare this with the right-hand side of (34). To do so, let us write that expression as $y(y^{j-1}x^{j-1})(s)$. Since $j < n$, and n is chosen as in (28), we have $d(x^{j-1}(s)) = j-1$, hence $d(y^{j-1}x^{j-1}(s)) = 0$. So what we need to show is that y, when applied to the depth-zero element $y^{j-1}x^{j-1}(s)$, gives the same output as when applied to the depth-zero element s. I claim that this will again hold by (28). To see that, we have to know how the element $y^{j-1}x^{j-1}(s)$, and in particular, its depth, behave under n successive applications of x. Now by (31), the first $j-1$ of these n applications of x simply strip away the same number of y's, increasing d by 1 at each step. Hence the result of applying x^{j-1} to $y^{j-1}x^{j-1}(s)$ is $x^{j-1}(s)$; so from that point on, we get the same outputs as when we apply the corresponding powers of x to s. So since (28) applies to the evaluation of $y(s)$, it also applies to the evaluation of $y(y^{j-1}x^{j-1}(s))$, and the resulting values are the same, as desired.

Finally, suppose we are in the case where $d(x^n(s)) = n$ for all n. Then by (30), $y(s)$ will be a y-stable element, i.e., will satisfy (25). This implies that it is fixed under $y^j x^j$, so the result of applying the left-hand side of (33) to s is $y(s)$. When we apply the right-hand side of (33), $y^j x^{j-1}$, to s, if we write that operation as $y(y^{j-1}x^{j-1})$, then the factor $y^{j-1}x^{j-1}$ will take s to an element s' again having $d(s') = 0$ and, by repeated application of (31), again satisfying $d(x^n(s')) = n$ for all n, and which, moreover, is $\approx$-equivalent to s, since they have the same images under x^{j-1}. So by the last condition of (30), the images of s and s' under y are the same. So the two sides of (33), applied to s, give the common value $y(s) = y(s')$, completing the proof of the theorem. □

Pace Nielsen has pointed out that the proof of Lemma 8 can be modified so that the only relations on x and y called on are $xyx = x$ and $x^{n-1}y^n x^n = yx^n$ for all $n > 0$ (the $(1,1,1)$ and $(n-1,n,n)$ cases of (2)). Namely, if we think of $x^{n-1}y^n x^n = yx^n$ as saying that $x^{n-1}y^n$ and y have the same effect on elements of $x^n S$, then this applies to elements $s \in \bigcap_{n\geq 0} x^n(S)$, and brings us directly to the relation $x^{n-1}y^n s = ys$ used in the above proof; after which only the relation $xyx = x$ is used. (Contrast this with the second assertion of Proposition 10, which shows that we cannot get such a result using only the family of relations $x^n y^n x^n = x^n$ and $y^n x^n y^n = y^n$.) Consequently, Theorem 11 can be strengthened to assert the equivalence of three conditions on an endomap x of a set: (i) condition (18), (ii) the existence of a strong inner inverse to x, and (iii) the existence of an element y satisfying the relations $xyx = x$, and $x^{n-1}y^n x^n = yx^n$ for all $n > 0$. The relations of (iii) are strictly weaker than y itself being a strong inner inverse to x, as may be seen from the construction, in §9 below, of the element y' of (50), which by (52) satisfies those relations, but by (51) does not satisfy $y' = y'xy'$.

It is natural to ask whether (18) is also a sufficient condition for an endomorphism x of a vector space V to have a strong inner inverse in $\mathrm{End}(V)$. Nielsen (personal communication) has an example showing that it is not.

8. A closer look at $k\mathcal{F}$

In this and the next section we shall, for conceptual simplicity, assume our division ring D is a field k, so that actions of $D\mathcal{F}$ on D-vector spaces V, which for general D do not have the property that the typical element acts by a D-vector-space endomorphism, become actions of $k\mathcal{F}$ by maps that are k-vector-space endomorphisms. But the reader will see that virtually nothing about k is used, so that the corresponding statements for $D\mathcal{F}$, if desired, are available.

So let k be a field and V a countable-dimensional k-vector space with basis $\{b_{n,h} \mid 1 \leq h \leq n\}$, and let x and y be the endomorphisms of V defined by

$$\begin{aligned} x(b_{n,h}) &= b_{n,h+1} \quad \text{if } h < n, \quad \text{while} \quad x(b_{n,n}) = 0, \\ y(b_{n,h}) &= b_{n,h-1} \quad \text{if } h > 1, \quad \text{while} \quad y(b_{n,1}) = 0. \end{aligned} \tag{35}$$

By Theorem 5, the monomials on the left-hand side of (2) form a k-basis of the subalgebra of $\mathrm{End}(V)$ generated by x and y, and so, in particular, are k-linearly independent.

But letting V_n denote the subspace of V spanned by $b_{n,1}, \dots, b_{n,n}$, consider the following four monomials in x and y, and their actions on V_n:

$$\begin{aligned} &1 \text{ fixes all } b_{n,h}, \\ &xy \text{ annihilates } b_{n,1}, \text{ and fixes all other } b_{n,h}, \\ &yx \text{ annihilates } b_{n,n}, \text{ and fixes all other } b_{n,h}, \\ &xyyx \text{ annihilates } b_{n,1} \text{ and } b_{n,n}, \text{ and fixes all other } b_{n,h}. \end{aligned} \tag{36}$$

These descriptions suggest that $1 + xyyx = xy + yx$ on each V_n.

What is wrong here?

The relation $1+xyyx = xy+yx$ does in fact hold on all V_n with $n > 1$. But for $n = 1$, the considerations suggesting that relation implicitly "double-count" the basis element $b_{1,1}$ in describing the effect of $xyyx$. On V_1, the elements xy, yx and $xyyx$ all act as 0, while 1 does not; so the asserted relation fails on V_1 – and only there.

This suggests that operators on V that, like $1 + xyyx - xy - yx$, have "small" images, are of interest in understanding $k\mathcal{F}$. Let us define on V, for each $i \geq 1$, operators ℓ_i and r_i, by

$$\begin{aligned} &\ell_i \text{ fixes, for each } n \geq i, \text{ the element } b_{n,i}, \text{ and annihilates all other} \\ &\quad \text{elements } b_{n,h} \ \ (n \geq 1,\ 1 \leq h \leq n), \\ &r_i \text{ fixes, for each } n \geq i, \text{ the element } b_{n,n+1-i}, \text{ and annihilates all other} \\ &\quad \text{elements } b_{n,h} \ \ (n \geq 1,\ 1 \leq h \leq n). \end{aligned} \tag{37}$$

Here ℓ and r are mnemonic for "left" and "right", since if we list the basis of V_n as $b_{n,1}, \dots, b_{n,n}$, then ℓ_i projects to the i-th basis element from the left (if any), and r_i to the i-th from the right (if any).

These operators are represented by elements of $k\mathcal{F}$. Namely, identifying elements of that algebra with their actions on V, it is not hard to check that for all $i \geq 1$,

$$\begin{aligned} \ell_i &= x^{i-1}y^{i-1} - x^i y^i, \\ r_i &= y^{i-1}x^{i-1} - y^i x^i. \end{aligned} \tag{38}$$

Elements with still "smaller" images are given by products of the above operators. Namely, for each $i, j \geq 1$,

$$\ell_i r_j \text{ fixes the single basis element } b_{i+j-1,i}, \text{ and annihilates all the other } b_{n,h}. \tag{39}$$

Let us note some relations which x, y, and the ℓ_i and r_i satisfy. First, the $i = 1$ cases of (38) give the following formulas, which can be applied to reduce any string of x's and y's which contains both letters to a linear combination of shorter strings:

$$xy = 1 - \ell_1, \quad yx = 1 - r_1. \tag{40}$$

In the next list of relations, (41), we make the temporary convention that ℓ_0 and r_0 represent the zero operator. Then we have the following equalities for all $i \geq 1$ (in which the $i = 1$ cases each say that a certain product is 0). These relations are easily deduced from (35) and (37), and allow us to eliminate all occurrences of ℓ's and r's to the left of x's and y's, and also all products $x\,r_1$ and $y\,\ell_1$.

$$\ell_i\, x = x\, \ell_{i-1}, \qquad r_{i-1}\, x = x\, r_i, \qquad \ell_{i-1}\, y = y\, \ell_i, \qquad r_i\, y = y\, r_{i-1}. \tag{41}$$

The $i = 1$ cases of the second and third relations above can be seen to generalize to

$$x^i r_i = 0, \quad y^i \ell_i = 0 \quad (i \geq 1). \tag{42}$$

Finally, from (37) it is easy to see that

$$\begin{aligned} &\ell_i^2 = \ell_i, \quad \ell_i \ell_j = 0 \text{ for } i \neq j, \\ &r_i^2 = r_i, \quad r_i r_j = 0 \text{ for } i \neq j, \\ &\ell_i r_j = r_j \ell_i \text{ for all } i, j. \end{aligned} \tag{43}$$

Using (40)-(43), we can reduce any element of $k\mathcal{F}$ to a k-linear combination of elements of the following four sorts.

(44) $\quad y^m \ \ (m \geq 1), \quad 1, \quad x^m \ \ (m \geq 1).$

(45) $\quad y^m \ell_i \ \ (i > m \geq 1), \quad \ell_i \ \ (i \geq 1), \quad x^m \ell_i \ \ (i, m \geq 1).$

(46) $\quad y^m r_i \ \ (i, m \geq 1), \quad r_i \ \ (i \geq 1), \quad x^m r_i \ \ (i > m \geq 1).$

(47) $\quad y^m r_i \ell_j \ \ (i, m \geq 1;\ j > m), \quad r_i \ell_j \ \ (i, j \geq 1), \quad x^m r_i \ell_j \ \ (j, m \geq 1;\ i > m).$

In fact, we have

Lemma 12. *The elements listed in* (44)-(47) *form a k-basis of $k\mathcal{F}$.*

Proof. We have seen that the elements (44)-(47) span $k\mathcal{F}$, so it will suffice to show that every nontrivial k-linear combination f of those elements has nonzero action on V.

Suppose first that f involves at least one of the elements in (44), say $u \in \{y^m, 1, x^m\}$, with nonzero coefficient in k. I claim that we can find an element $b_{n,i}$ in our basis for V which is annihilated by all the monomials occurring with nonzero coefficient in f that lie in (45)-(47), but not by u. Indeed, for each $n > 0$, each term in (45)-(47) has nonzero action on at most one of $b_{n,1}, \ldots, b_{n,n}$, while the number of members of each such set annihilated by our element u is bounded, independent of n, by the exponent on x or y in u (if any). Hence for large enough n, there exists a $b_{n,i}$ which is neither an element on which the finitely many terms of f in (45)-(47) have nonzero value, nor one annihilated by u; let us choose any such $b_{n,i}$. Note that each member of (44) other than u which is nonzero on $b_{n,i}$ carries it to a basis element other than $u(b_{n,i})$, since distinct members of (44) shift second subscripts of the $b_{n,i}$ by different amounts (35). Hence in $f(b_{n,i})$, the basis-element $u(b_{n,i})$ has nonzero coefficient; so f has nonzero action on V.

Suppose, next, that f involves no members of (44), but has some member of (45) or (46) with nonzero coefficient. Assume without loss of generality that it involves an element u of (45), and let ℓ_i be the ℓ-factor occurring. Thus, u acts nontrivially only on basis elements $b_{n,i}$ $(n \geq i)$, and I claim we can find some n for which u, but none of the elements of (46) or (47) occurring in f, does so. Note that u annihilates $b_{n,i}$ for at most finitely many n. Each element of (46) acts nontrivially on members of our basis that are a fixed distance from the *right* end of the families $\{b_{n,1}, \ldots, b_{n,n}\}$, so for n large enough, none of the finitely many such elements which occur in f will act nontrivially on the i-th element from the *left*. Moreover, each member of (47) is nonzero on $b_{n,i}$ for at most one n. Hence for all but finitely many n, the element $b_{n,i}$ has nonzero image under u but not under any of the members of (46) or (47) occurring in f; let us pick such a $b_{n,i}$. Of the other members of (45) occurring in f, those that involve ℓ_j for some $j \neq i$ annihilate $b_{n,i}$, while those that involve ℓ_i but begin with a different factor x^m, 1 or y^m send $b_{n,i}$ to a different basis element. So again, the basis element $u(b_{n,i})$ has nonzero coefficient in $f(b_{n,i})$, so f has nonzero action on V.

Finally, if f involves no members of (44)-(46), we take a member u of (47) that it involves, and let $r_i\ell_j$ be the factors other than a power of x or y in that operator. Then $u(b_{i+j-1,j}) \neq 0$, and we easily check that the other elements of (47) appearing in f either annihilate $b_{i+j-1,j}$, or send it to a basis element different from the one to which u sends it; so $f(b_{i+j-1,j}) \neq 0$. □

Although $\mathcal{F}$ has trivial center, $k\mathcal{F}$ has nontrivial central idempotents. Namely, for each $n \geq 1$, the element

(48) $\quad p_n \ = \ r_n\ell_1 + r_{n-1}\ell_2 + \cdots + r_1\ell_n$

acts on V by projection to V_n; and since all members of $k\mathcal{F}$ carry V_n into itself, p_n commutes with all such elements. (For results on the centers of monoid algebras of free inverse monoids on more than one generator, see [9].)

The above material on the structure of $k\mathcal{F}$ has considerable overlap with [1, §4]. In particular, our $\ell_i r_j$ and p_n would be, in the notation of display (4.4) and Lemma 4.3 of that note, $q_{-i+1,j-1}$, and h_{n-1}; and Proposition 4.5 of that paper shows, inter alia, that the latter elements generate the socle of $k\mathcal{F}$ (which is equivalent to saying that the socle is spanned over k by the former elements).

Though in the above arguments, we have been regarding $k\mathcal{F}$ as an algebra of operators on the vector space $V = \bigoplus V_n$, the results on its structure that we have deduced necessarily hold for the abstract algebra

$k\mathcal{F}$; so we can look at the features discussed above in connection with any action of $k\mathcal{F}$ on any vector space. For instance, if x is an *invertible* endomorphism of a vector space V, and y its inverse, then we get an action in which, it is easy to see, all ℓ_i and r_i act trivially (i.e., as the zero operator). If x acts by a one-to-one but not invertible endomorphism of a space V, and we let y act by a left inverse thereof, we get an action under which the r_i are trivial, but not the ℓ_i; and dually if x is surjective but not one-to-one. In these cases, all the products $r_i\,\ell_j$, and hence the central idempotents p_n, act trivially.

Note that there is a homomorphism $\mathcal{F} \to \mathbb{Z}$ taking x to 1 and y to -1, and that this induces a grading on $k\mathcal{F}$, under which the basis elements (44)-(47) are homogeneous: those whose expressions begin with y^m have degree $-m$, those beginning with x^m have degree $+m$, and those beginning with neither have degree 0. The homogeneous component of $k\mathcal{F}$ of degree 0 is commutative, generated over k by the idempotent elements ℓ_i and r_i.

The following result, which relates the behavior of r_i and ℓ_i with that of x under an arbitrary action of $k\mathcal{F}$ on a k-vector space, will be used in the next section.

Lemma 13. *Let V be a k-vector space given with a left action of $k\mathcal{F}$, and i a positive integer. Then for $i \geq 1$,*

(i) *The action of the projection map r_i on V annihilates the subspace* $\ker(x^{i-1})$, *and has as image a complement of that subspace in the (generally larger) subspace* $\ker(x^i)$.

(ii) *The action of the projection map ℓ_i on V annihilates the subspace* $\mathrm{im}(x^i)$, *and has as image a complement of that subspace in the (generally larger) subspace* $\mathrm{im}(x^{i-1})$.

(*In these statements, x^0 is understood to be the identity operator. Thus, in the $i = 1$ case of* (i), *since* $\ker(\mathrm{id}) = 0$, *the conclusion simply says that the projection r_1 has image* $\ker(x)$; *and in* (ii), *since* $\mathrm{im}(\mathrm{id}) = V$, *the conclusion says that ℓ_1 is a projection of V along* $\mathrm{im}(x)$ *onto a complement of that subspace.*)

The corresponding statements hold with x replaced by y, and the roles of ℓ_i and r_i interchanged.

Proof. Since r_i and ℓ_i are idempotent elements of $k\mathcal{F}$, they act on V by projection operators.

Since $r_i = y^{i-1}x^{i-1} - y^ix^i$, this element is right divisible by x^{i-1}, hence annihilates $\ker(x^{i-1})$; while its image lies in $\ker(x^i)$ by the first equation of (42). Since it is a projection, to show that its image is a complement of $\ker(x^{i-1})$ in $\ker(x^i)$, it remains only to show that the only elements of $\ker(x^i)$ annihilated by r_i are the elements of $\ker(x^{i-1})$. And indeed, if $v \in \ker(x^i)$ is annihilated by $r_i = y^{i-1}x^{i-1} - y^ix^i$, then it is annihilated by $y^{i-1}x^{i-1}$, hence by that element's left multiple $x^{i-1}y^{i-1}x^{i-1} = x^{i-1}$, as required.

Turning to ℓ_i, this annihilates $\mathrm{im}(x^i)$ by repeated application of the first equation of (41). Since $\ell_i = x^{i-1}y^{i-1} - x^iy^i$, it is left divisible by x^{i-1}, so its image lies in $\mathrm{im}(x^{i-1})$. Thus, as above, it remains to show that any element of $\mathrm{im}(x^{i-1})$ annihilated by ℓ_i lies in $\mathrm{im}(x^i)$. And indeed, the condition that an element $x^{i-1}(v) \in \mathrm{im}(x^{i-1})$ be annihilated by ℓ_i is $x^{i-1}y^{i-1}x^{i-1}(v) - x^iy^ix^{i-1}(v) = 0$. Applying (2) to each term of this relation, it becomes $x^{i-1}(v) - x^iy(v) = 0$, showing that $x^{i-1}(v)$ indeed lies in $\mathrm{im}(x^i)$.

The final assertion is clear by symmetry. (We will not need it below.) □

9. Inner inverses satisfying other systems of relations

In this section, we regard the monoid relations (2), (3), (4) by which we defined $\mathcal{F}$ as k-algebra relations defining $k\mathcal{F}$.

It is natural to ask whether the relations defining $k\mathcal{F}$ comprise the "strongest" family of k-algebra relations that one can force an inner inverse y of a general element x of a "good" regular k-algebra (such as an infinite product of full matrix algebras over k) to have.

An easy observation which suggests a negative answer is the following: It is known that for all n the ring of $n \times n$ matrices over k is *unit regular,* i.e., that for every x there exists an *invertible* y such that $xyx = x$ [7, Lemma 1.3]. Now if x is not invertible and y is, we can't have $yxy = y$; so for noninvertible x, the condition that y be invertible is incompatible with the relations (2). Unfortunately, the condition that y be invertible is not a k-algebra relation in x and y; so the above does not contradict the possibility that the relations defining $k\mathcal{F}$ might be the strongest set of k-algebra relations *in those two elements* that we can force an inner inverse y to have in the above ring.

We shall see at the end of this section that a modification of the above idea does show that the set of relations defining $k\mathcal{F}$ is not a greatest element of the partially ordered set of such families of relations. But let us first show that it *is* a *maximal* element of that partially ordered set. For this, we will need

Corollary 14 (to Lemma 13)**.** *An action of $k\mathcal{F}$ on a vector space V is faithful if and only if for all positive integers i,*

$$\text{(49)} \qquad \mathrm{im}(x^i) \cap \ker(x) \;\neq\; \mathrm{im}(x^{i-1}) \cap \ker(x).$$

Proof. Clearly, the left-hand side of (49) is contained in the right-hand side; so it suffices to show that the given action is faithful if and only if for every i, there is an element of the right-hand side of (49) that is not in the left-hand side.

It is easy to verify that every nonzero two-sided ideal of $k\mathcal{F}$ contains at least one of the elements (47); and with the help of (41), one can deduce further that every such ideal contains an element $\ell_i r_1 \;\; (i \geq 1)$. We shall now show that for each i, the existence of an element belonging to the right-hand side of (49) but not the left is equivalent to the condition that $\ell_i r_1$ act nontrivially on V, i.e., not be in the kernel of our action. Thus, the inequalities (49) will hold for all i if and only if that kernel is zero, giving our desired result.

Suppose first that some $v \in V$ belongs to $\mathrm{im}(x^{i-1}) \cap \ker(x)$ but not to $\mathrm{im}(x^i) \cap \ker(x)$. Since $v \in \ker(x)$, we see from the parenthetical statement in Lemma 13 that it is fixed under r_1, while since it lies in $\mathrm{im}(x^{i-1})$ but not in $\mathrm{im}(x^i)$, statement (ii) of that lemma shows that it is not annihilated by ℓ_i. Hence $\ell_i r_1(v) \neq 0$, so $\ell_i r_1$ indeed acts nontrivially.

Conversely, suppose $\ell_i r_1$ acts nontrivially on some $v \in V$. Then I claim that $\ell_i r_1(v)$ lies in the right-hand side of (49) but not in the left-hand side. Indeed, since that element is a nonzero element of $\ell_i(V)$, Lemma 13(ii) tells us that it belongs to $\mathrm{im}(x^{i-1})$ but not to $\mathrm{im}(x^i)$, while since $\ell_i r_1 = r_1 \ell_i$, it also lies in $r_1(V) = \ker(x)$, as required. □

We deduce

Proposition 15. *Let R be a k-algebra, and φ and φ' k-algebra homomorphisms $k\mathcal{F} \to R$ such that $\varphi(x) = \varphi'(x)$. Then φ is one-to-one if and only if φ' is.*

Hence no proper *homomorphic image $(k\mathcal{F})/I$ has the property that for every element x in a direct product R of full matrix rings over k, there exists $y \in R$ which, with x, satisfies the relations of $(k\mathcal{F})/I$.*

Thus, the set of k-algebra relations satisfied by x and y in $k\mathcal{F}$ is maximal among sets S of k-algebra relations in two noncommuting indeterminates such that for every element x in a direct product R of full matrix algebras, one can find a $y \in R$ such that x and y together satisfy S.

Proof. Embedding R in an algebra of the form $\mathrm{End}(V)$, we get from the preceding corollary a condition which is necessary and sufficient both for φ and for φ' to be one-to-one, in terms of the action of x on V. Since $\varphi(x) = \varphi'(x)$, this yields the first assertion.

We know from the second paragraph of Theorem 5 that there exist elements x and y in an infinite direct product of matrix rings over k which satisfy the relations of $k\mathcal{F}$ but no others, hence the preceding assertion shows that for such x, there exists no y' such that x and y' satisfy the relations of a proper homomorphic image $(k\mathcal{F})/I$ of $k\mathcal{F}$.

The third assertion then follows. □

One can easily strengthen the above proof to show that for φ and φ' as in the first sentence of the proposition, their kernels have the same intersection with the ideal of $k\mathcal{F}$ spanned by the elements (47) (the socle of $k\mathcal{F}$). It seems likely that these kernels must in fact be equal; I leave this for others to investigate.

Let us now prove, on the other hand, that the set of relations satisfied by x and y in $k\mathcal{F}$ is not the only maximal set of k-algebra relations that can be forced in this way. We will use the idea of the second paragraph of this section in our example, but instead of trying to make y everywhere invertible, we shall only do this on x-basic subspaces of one chosen dimension m. To see how, let us again think in terms the action of $k\mathcal{F}$ on the vector space $V = \bigoplus V_n$ spanned by elements $b_{n,h}$, as in §8. If we let

$$\text{(50)} \qquad y' \;=\; y \,+\, x^{m-1}\ell_1 r_m,$$

we see that y' acts like y on the spaces V_n with $n \neq m$ (since $\ell_1 r_m$ acts trivially on those spaces), but that on V_m, it cyclically permutes the m basis elements $b_{m,i}$. Since x does not act invertibly on V_m, neither does $y'xy'$, so since y' does, we have

$$\text{(51)} \qquad y' \;\neq\; y'x\,y'.$$

On the other hand, the reader can easily verify, by calculation in $\mathrm{End}(V)$, that for all natural numbers i,

$$\text{(52)} \qquad (y')^i\,x^i \;=\; y^i\,x^i, \quad \text{and} \quad x^i\,(y')^i \;=\; x^i\,y^i.$$

Hence the subalgebra of $k\mathcal{F}$ generated by x and y' contains all ℓ_i and r_i, and these are expressible in terms of x and y' by the same formulas (38) that express them in terms of x and y. In particular, that subalgebra contains $y = y' - x^{m-1}\ell_1 r_m$ (see (50)), hence it is all of $k\mathcal{F}$. From the $i = 1$ case of (52), we also see that $x = xy'x$; i.e., y' is an inner inverse to x.

The contrast between y and y' can now be attested concretely by the easily checked relations

$$(y')^m p_m \;=\; p_m, \qquad y^m p_m = 0. \tag{53}$$

Since ℓ_i and r_i can be expressed in terms of x and y' by the same formulas that express them in terms of x and y, these can be regarded as relations in $k\mathcal{F}$ satisfied by x and y' and by x and y, respectively.

Thus, we get

Proposition 16. *In the k-algebra $k\mathcal{F}$, the element y' given by* (50) *is another inner inverse to x, which together with x generates the whole algebra, but which, by* (53)*, satisfies with x a family of relations incomparable with the family of relations satisfied by y. Hence these two sets of relations are distinct maximal elements among the families S of k-algebra relations referred to in the last sentence of Proposition 15.* □

This leaves open the question of whether every algebra presented by generators x and y, and a maximal set of relations which include the relation $x = xyx$ and which can be "forced" in the sense we have been discussing, must be isomorphic to $k\mathcal{F}$ by an isomorphism fixing x. In addition to questions about relations satisfied by x and y alone, one might want to look at variants, such as what relations can be forced on an *invertible* inner inverse y to an element x.

10. Acknowledgements

I am indebted to Pace Nielsen, Kevin O'Meara and Janez Šter for stimulating interchanges leading to this material, to Ben Steinberg and Mark Lawson for information on inverse monoids, and to Warren Dicks, Ken Goodearl, and T. Y. Lam for further discussion.

References

[1] P. Ara and K. R. Goodearl, *The realization problem for some wild monoids and the Atiyah problem,* 48 pp., http://arXiv.org/abs/1410.8838v2 . To appear, Transactions of the A. M. S..

[2] Giuseppe Baccella, *Semiartinian V-rings and semiartinian von Neumann regular rings,* J. Algebra **173** (1995) 587–612. http://dx.doi.org/10.1006/jabr.1995.1104 . MR1327870

[3] George M. Bergman, *The diamond lemma for ring theory,* Advances in Mathematics **29** (1978) 178–218. MR0506890.

[4] George M. Bergman, *An Invitation to General Algebra and Universal Constructions,* 2015, Springer Universitext, x+572 pp.. http://dx.doi.org/10.1007/978-3-319-11478-1 . MR3309721

[5] A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups. Vol. I.,* AMS Mathematical Surveys, No. 7, 1961, xv+224 pp.. http://www.ams.org/books/surv/007.1/ . MR0132791

[6] A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups. Vol. II.,* AMS Mathematical Surveys, No. 7, 1967, xv+350 pp.. http://www.ams.org/books/surv/007.2/ . MR0218472

[7] K. R. Goodearl, *von Neumann regular rings*, second edition. Robert E. Krieger Publishing Co., Inc., Malabar, FL, 1991. xviii+412 pp. MR1150975. (First ed.: MR0533669)

[8] Mark V. Lawson, *Inverse semigroups. The theory of partial symmetries,* World Scientific Publishing Co., Inc., 1998. xiv+411 pp.. MR1694900.

[9] M. J. Crabb and W. D. Munn, *The centre of the algebra of a free inverse monoid,* Semigroup Forum **55** (1997) 215–220. MR1457767

[10] T. Y. Lam, *A first course in noncommutative rings,* Second edition. Springer Graduate Texts in Mathematics, 131. 2001. xx+385 pp. MR1838439

[11] W. D. Munn, *Inverse semigroup rings,* Topics in algebra, Part 1 (Warsaw, 1988), 65–73, Banach Center Publ., 26, Part 1, PWN, Warsaw, 1990. MR1171226

[12] W. D. Munn, *Free inverse semigroups,* Proc. London Math. Soc. (3) **29** (1974) 385–404. MR0360881

[13] Pace P. Nielsen and Janez Šter, *Connections between unit-regularity, regularity, cleanness, and strong cleanness of elements and rings.* To appear, Transactions of the A. M. S..

[14] Mario Petrich, *Inverse semigroups,* John Wiley & Sons, Pure and Applied Mathematics, 1984. x+674 pp.. MR0752899.

University of California, Berkeley, CA 94720-3840, USA
E-mail address: gbergmammath.berkeley.edu